

Special Topics in Operations Research 16:711:611

Convex Analysis and Optimization

Spring 2009

Rutgers University

Prof. Eckstein

Solutions to Homework 3

1. (a) Consider any $y = (y^1, \dots, y^m) \in C_1^* \times \dots \times C_m^*$. Then for any $x = (x^1, \dots, x^m) \in C_1 \times \dots \times C_m$, we have $\langle x^i, y^i \rangle \leq 0$ for $i = 1, \dots, m$, since $x^i \in C_i$ and $y^i \in C_i^*$. Thus,

$$\langle x, y \rangle = \sum_{i=1}^m \langle x^i, y^i \rangle \leq 0,$$

because each term in the sum is nonpositive. Since $x \in C_1 \times \dots \times C_m$ was arbitrary, $y \in (C_1 \times \dots \times C_m)^*$. Since $y \in C_1^* \times \dots \times C_m^*$ was arbitrary, $C_1^* \times \dots \times C_m^* \subseteq (C_1 \times \dots \times C_m)^*$.

For the reverse inclusion, consider any $y = (y^1, \dots, y^m) \notin C_1^* \times \dots \times C_m^*$. This means that for at least one $i \in \{1, \dots, m\}$, we have $y^i \notin C_i^*$. This in turn means that there exists $x^i \in C_i$ such that $\langle x^i, y^i \rangle > 0$. Consider now the vector $x = (0, \dots, 0, x^i, 0, \dots, 0)$, where the nonzero entry is in the i^{th} position. Since the C_j , $j \neq i$, are cones, they contain 0, and so $x \in C_1 \times \dots \times C_m$. But then we have $\langle x, y \rangle = \langle x^i, y^i \rangle > 0$, so $y \notin (C_1 \times \dots \times C_m)^*$. Thus, in view of the above inclusion, $C_1^* \times \dots \times C_m^* = (C_1 \times \dots \times C_m)^*$.

- (b) Consider any $y \in \bigcap_{i \in I} C_i^*$. For any $x \in \bigcup_{i \in I} C_i$, there must exist $j \in I$ such that $x \in C_j$. Since we must have $y \in C_j^*$, we have $\langle x, y \rangle \leq 0$. This logic establishes that $\bigcap_{i \in I} C_i^* \subseteq (\bigcup_{i \in I} C_i)^*$. Now consider any $y \notin \bigcap_{i \in I} C_i^*$. Then there exists $j \in I$ such that $y \notin C_j^*$, which in turn means there exists $x \in C_j$ such that $\langle x, y \rangle > 0$. Since this x is in $\bigcup_{i \in I} C_i$, it follows that $y \notin (\bigcup_{i \in I} C_i)^*$. It then follows that $\bigcap_{i \in I} C_i^* = (\bigcup_{i \in I} C_i)^*$.
- (c) Consider $y \in C_1^* \cap C_2^*$, and any $x \in C_1 + C_2$. Now, we must have $x = x^1 + x^2$, where $x^1 \in C_1$ and $x^2 \in C_2$. Since $y \in C_1^*$, we have $\langle x^1, y \rangle \leq 0$, and since $y \in C_2^*$, we also have $\langle x^2, y \rangle \leq 0$. Therefore, $\langle x, y \rangle = \langle x^1 + x^2, y \rangle = \langle x^1, y \rangle + \langle x^2, y \rangle \leq 0 + 0 = 0$, and $y \in (C_1 + C_2)^*$. Thus, $C_1^* \cap C_2^* \subseteq (C_1 + C_2)^*$. Next, consider $y \notin C_1^* \cap C_2^*$. Then for either $i = 1$ or $i = 2$, we have $y \notin C_i^*$. There must then exist some $x \in C_i$ with $\langle x, y \rangle > 0$. But this x is also in $C_1 + C_2$ (just add the vector 0 from the other cone), and so $y \notin (C_1 + C_2)^*$. So, $C_1^* \cap C_2^* = (C_1 + C_2)^*$.

2. There is a nearly trivial proof based on the polar cone theorem: if there does not exist an $a \in K^*$ with $\langle a, z \rangle > 0$, then $\langle y, z \rangle \leq 0$ for all $y \in K^*$, which means that $y \in K^{**}$. Since K is nonempty and closed, the polar cone theorem asserts that $K^{**} = K$, so $z \in K$, contradicting the hypothesis.

There are several alternative approaches using more basic principles. Here is one: since K is a closed convex set, there exists some $a \in \mathbb{R}^n \setminus \{0\}$ and $b \in \mathbb{R}$ such that

$\langle a, z \rangle > b$ and $\langle a, x \rangle \leq b$ for all $x \in K$. Since $0 \in K$, we must have $0 = \langle a, 0 \rangle \leq b$, and thus $\langle a, z \rangle > 0$. Furthermore, if there were any $x \in K$ such that $\langle a, x \rangle > 0$, then for sufficiently large $\alpha > 0$ we would have $\alpha x \in K$ and $\langle a, \alpha x \rangle = \alpha \langle a, x \rangle > b$, a contradiction. So we know $\langle a, x \rangle \leq 0$ for all $x \in K$, that is, $a \in K^*$.

3. We start by reasoning similarly to part (c) above. Take any $y \in C_1^* + C_2^*$. Then $y = y^1 + y^2$, where $y^1 \in C_1$ and $y^2 \in C_2$. Now consider any $x \in C_1 \cap C_2$. Since $x \in C_1$ and $y^1 \in C_1^*$, we have $\langle x, y^1 \rangle \leq 0$. Since we also have $x \in C_2$, we similarly have $\langle x, y^2 \rangle \leq 0$. We then calculate $\langle x, y \rangle = \langle x, y^1 + y^2 \rangle = \langle x, y^1 \rangle + \langle x, y^2 \rangle \leq 0 + 0 = 0$. We then conclude that $C_1^* + C_2^* \subseteq (C_1 \cap C_2)^*$. In general, we may not know that $C_1^* + C_2^*$ is closed, but since $(C_1 \cap C_2)^*$ must be a closed set and contains $C_1^* + C_2^*$, it certainly also contains $\text{cl}(C_1^* + C_2^*)$.

For the opposite inclusion, consider some $z \notin \text{cl}(C_1^* + C_2^*)$. Because $\text{cl}(C_1^* + C_2^*)$ is a closed convex cone, we may invoke problem 2 to conclude that there exists some $a \in [\text{cl}(C_1^* + C_2^*)]^*$ such that $\langle a, z \rangle > 0$. Now, as shown in class, $(\text{cl } X)^* = X^*$ for any set X , so

$$\begin{aligned} a &\in (C_1^* + C_2^*)^* \\ &= C_1^{**} \cap C_2^{**} \quad [\text{by problem 1(c)}] \\ &= C_1 \cap C_2 \quad [\text{by polar cone theorem, since } C_1, C_2 \text{ are closed convex cones}]. \end{aligned}$$

Since $\langle a, z \rangle > 0$ and $a \in C_1 \cap C_2$, $z \notin (C_1 \cap C_2)^*$. We conclude that $(C_1 \cap C_2)^* = \text{cl}(C_1^* + C_2^*)$.

4. (a) Suppose $\{x^k\} \subseteq K$ is a convergent sequence whose limit is x . Then $Ax^k \rightarrow Ax$, and since $Ax^k \in C$ for all k and C is closed, we have $Ax \in C$. Therefore, $x \in K$, and we can deduce that K is a closed set.

Take any $x \in K$ and $\alpha \geq 0$. Then $A(\alpha x) = \alpha \cdot (Ax) \in C$, because $Ax \in C$ and C is a cone. This establishes that K is a cone.

Finally, take any $x^1, x^2 \in K$. Then $A(x^1 + x^2) = Ax^1 + Ax^2 \in C$ because $Ax^1, Ax^2 \in C$ and C is a convex cone. Since we already established that K is a cone, it must be convex.

- (b) Suppose $z \in P$ and $\alpha \geq 0$. Then $z = A^\top y$ for some $y \in C^*$; since C^* is a cone, $\alpha y \in C^*$ and thus $A^\top(\alpha y) = \alpha A^\top y = \alpha z \in P$. This establishes that P is cone. It follows that $\text{cl } P$ is also a cone (I omit the proof, but it is very simple).

Consider any $x \in K$ and $z \in P$. Then $z = A^\top y$ for some $y \in C^*$, and $\langle x, z \rangle = \langle x, A^\top y \rangle = \langle Ax, y \rangle \leq 0$, the last inequality following from $Ax \in C$ and $y \in C^*$. This establishes that $P \subseteq K^*$. Since K^* must be a closed set and contains P , we also have $\text{cl } P \subseteq K^*$.

For the reverse inclusion, consider any $z \notin \text{cl } P$. From problem 2, there must exist $q \in (\text{cl } P)^* = P^*$ such that $\langle q, z \rangle > 0$. Since $q \in P^*$, we have $\langle q, w \rangle \leq 0$ for all $w \in P$, that is, $\langle q, A^\top y \rangle \leq 0$ for all $y \in C^*$. Equivalently, $\langle Aq, y \rangle \leq 0$ for all $y \in C^*$, which means $Aq \in C^{**} = C$ (because C is a closed convex cone). From the definition of K , this means $q \in K$, and since $\langle q, z \rangle > 0$, we therefore have $z \notin K^*$. Thus, we have $\text{cl } P = K^*$.

(c) We have just established $\text{cl } P = K^*$. We then have

$$P^* = (\text{cl } P)^* = K^{**} = K.$$

The justifications for the above three equalities are, respectively,

- $X^* = (\text{cl } X)^*$ for any set X
 - Taking the polar of both sides in $\text{cl } P = K^*$
 - The polar cone theorem, since K is closed and convex.
5. (a) We start by considering the case $n = 1$, in which case $K = [0, \infty)$. Then $K^* = \{y \in \mathbb{R} \mid xy \leq 0 \forall x \in [0, \infty)\} = (-\infty, 0] = -K$. For $n > 1$, we note that $K = [0, \infty)^n$, and so problem 1(a) implies that $K^* = ([0, \infty)^n)^* = ([0, \infty)^*)^n = (-\infty, 0]^n = -K$.
- (b) First, we note that

$$\begin{aligned} -K &= \{-(x, w) \in \mathbb{R}^n \times \mathbb{R} \mid w \geq \|x\|\} \\ &= \{(y, z) \in \mathbb{R}^n \times \mathbb{R} \mid -z \geq \|-y\|\} \\ &= \{(y, z) \in \mathbb{R}^n \times \mathbb{R} \mid z \leq -\|-y\|\} \\ &= \{(y, z) \in \mathbb{R}^n \times \mathbb{R} \mid z \leq -\|y\|\}. \end{aligned}$$

Now consider any $(x, w) \in K$ and $(y, z) \in -K$. Then

$$\begin{aligned} \langle (x, w), (y, z) \rangle &= \langle x, y \rangle + wz \\ &\leq \|x\| \|y\| + wz && \text{[by the Cauchy-Schwarz inequality]} \\ &\leq \|x\| \|y\| - \|x\| \|y\| && \text{[since } w \geq \|x\| \text{ and } z \leq -\|y\|\text{]} \\ &= 0. \end{aligned}$$

Thus, we conclude that $-K \subseteq K^*$. Now consider any $(u, v) \in (\mathbb{R}^n \times \mathbb{R}) \setminus (-K)$. Then we must have $v > -\|u\|$. We now distinguish two cases:

- $u = 0$: In this case, $v > 0$. Considering the vector $(0, 1) \in \mathbb{R}^n \times \mathbb{R}$, we have $\langle (0, 1), (u, v) \rangle = v > 0$. Since $(0, 1) \in K$, we conclude that $(u, v) \notin K^*$.
- $u \neq 0$: Consider the vector $(u, \|u\|) \in K$. Then

$$\begin{aligned} \langle (u, v), (u, \|u\|) \rangle &= \|u\|^2 + \|u\| \cdot v \\ &> \|u\|^2 - \|u\|^2 && \text{[because } v > -\|u\| \text{ and } \|u\| \neq 0\text{]} \\ &= 0. \end{aligned}$$

Since $(u, \|u\|) \in K$, we conclude that $(u, v) \notin K^*$.

Combining these two cases and that we have already proved $-K \subseteq K^*$, we must have $-K = K^*$.